arXiv:1603.02473v3 [math.NT] 15 Mar 2016

On the parity of the multiplicative order of certain products of integers related to Gauss factorials

Timothy Foo

Abstract

In this note, we prove that under some conditions, certain products of integers related to Gauss factorials are always quadratic residues.

Introduction

Let n be a positive integer and p a prime congruent to 1 modulo n. Then it follows that the numbers $\{1, 2, \ldots, p-1\}$ may be divided into $(p-1)/n$ equally sized sets. In analogy with Wilson's theorem, it is interesting to ask about the products $\prod_{j=(k-1)(p-1)/n+1}^{k(p-1)/n} j$, where $k = 1, \ldots, n$. The interesting topic of Gauss factorials was studied in [2-9], and in [5] the partial products to be studied here, which are related to Gauss factorials, were and henceforth here are denoted

$$\Pi_k^{(n)} = \prod_{j=(k-1)(p-1)/n+1}^{k(p-1)/n} j$$

where $k = 1, \ldots, n$. In particular, in [2-9], interest has been on the multiplicative order of Gauss factorials, and the accompanying conditions on p. In [14], Mordell proved that for $p \equiv 3 \bmod 4$,

$$\Pi_1^{(2)} \equiv (-1)^a \bmod p$$

where $a \equiv \frac{1}{2}(1 + h(-p)) \bmod 2$, $h(-p)$ being the class number of the imaginary quadratic field $\mathbb{Q}[\sqrt{-p}]$. It is interesting to ask, is there a statement for products over various k of $\Pi_k^{(q)}$, q an odd prime, that relates to the class number? When p is a prime congruent to 3 modulo 4, the multiplicative order of an element in $(\mathbb{Z}/p\mathbb{Z})^*$ is either odd or congruent to 2 modulo 4. The aim of this note is to show that for primes p congruent to 3 modulo 4, and also being congruent to 1 modulo q so that the appropriate products $\Pi_k^{(q)}$, where $k = 1, \ldots, q$, exist, certain products of these products have odd multiplicative order modulo p. For p congruent to 3 modulo 4, this is equivalent to these elements being quadratic residues modulo p. Thus, we have the following theorem.

Theorem 1. *Let q be an odd prime. Let p be a prime such that $p \equiv 3 \bmod 4$ and $p \equiv 1 \bmod q$. Then the element*

$$\prod_{\substack{k=1 \\ (q+1)/2-k \equiv 1 \bmod 2}}^{(q-1)/2} \Pi_k^{(q)}$$

is a quadratic residue modulo p.

In the remainder of this note, we shall prove Theorem 1 and discuss its implications. Then we look at further generalizations. Some notation is helpful which we introduce. For $a \in (\mathbb{Z}/p\mathbb{Z})^*$, let the Legendre symbol be denoted by

$$\left(\frac{a}{p}\right) = \begin{cases} 1, \text{ if a is a quadratic residue modulo p} \\ -1 \text{ otherwise.} \end{cases}$$

Furthermore, as stated earlier, let $h(-p)$ denote the class number of the imaginary quadratic field $\mathbb{Q}[\sqrt{-p}]$, $p \equiv 3 \bmod 4$. Finally, for $p \equiv 1 \bmod q$, let a_k and b_k denote respectively the number of quadratic residues and nonresidues in the set $\{(k-1)(p-1)/q+1, \ldots, k(p-1)/q\}$.
At this point, let us mention Theorem 3.1 of [13], an example of which is seen in [12]. Specifically, it follows from this theorem that if $q \equiv 3 \bmod 4$, H the subgroup of squares in $(\mathbb{Z}/q\mathbb{Z})^*$, $\beta = (\sum_{\substack{i=1 \\ -i \in H}}^{q-1} i)/q$, $p \equiv 1 \bmod q$, $4p^{h(-q)} = a^2 + qb^2$, where $a \equiv 2 \bmod q$, then

$$\left(\frac{a}{p}\right) = \left(\frac{(-1)^{\beta} \prod_{-i \in H} \left(i\left(\frac{p-1}{q}\right)\right)!}{p}\right). \tag{0.1}$$

We have that

Theorem 2. *Let p, q be primes, $q \equiv 3 \bmod 4$, $p \equiv 1 \bmod q$, $p \equiv 3 \bmod 4$. Let a be defined as above. Then the following are equivalent.*
(1)

Theorem 1 applied to the case when q is 3 mod 4 $\Leftrightarrow$ Equation (0.1)

(2)

$$\left(\frac{a}{p}\right) = (-1)^{(q+1)/4}.$$

Proof. Since $\beta = \frac{h(-q)+1}{2} + \frac{q-3}{4}$ and $q \equiv 3 \bmod 4$,

$$\begin{aligned}
1 &= \left(\frac{\prod_{\substack{k=1 \\ (q+1)/2-k \equiv 1 \bmod 2}}^{(q-1)/2} \Pi_k^{(q)}}{p}\right) \text{ by Theorem 1} \\
&= \left(\frac{\prod_{i=1}^{(q-1)/2} \left(i\left(\frac{p-1}{q}\right)\right)!}{p}\right) \\
&= (-1)^{1+(h(-q)+1)/2} \left(\frac{\prod_{-i \in H} \left(i\left(\frac{p-1}{q}\right)\right)!}{p}\right) \\
&= (-1)^{(q+1)/4} \left(\frac{a}{p}\right) \text{ by Equation (0.1).}
\end{aligned}$$

The expression for the class number of an imaginary quadratic field in terms of Legendre symbols

The proof of Theorem 1 here begins with the beautiful expression of the class number of the imaginary quadratic field $\mathbb{Q}[\sqrt{-p}]$, $p \equiv 3 \bmod 4$, due to Dirichlet. The formula is given by

$$h(-p) = \frac{1}{2 - \left(\frac{2}{p}\right)} \sum_{a=1}^{(p-1)/2} \left(\frac{a}{p}\right)$$

where $p \equiv 3 \bmod 4$, $p > 3$ and references include Corollary 5.3.13 of [1], and [10] (equations (7), (8) of Chapter 1 and equation (19) of chapter 6). This nice formula is due to the Legendre symbol being an odd character when $p \equiv 3 \bmod 4$. A different form of this formula, generalized to the other primes, instead of the prime 2, may be obtained by similar treatment of the L-function $L(1, (\frac{-}{p}))$ following the outline summarized in Chapter 1 of [10]. This becomes our first lemma.

Lemma 1. *Let q be an odd prime, and $p \equiv 3 \bmod 4$, $p > 3$. Then for $p \neq q$,*

$$h(-p) = \frac{1}{q - \left(\frac{q}{p}\right)} \sum_{a=1}^{(p-1)/2} \left(\frac{a}{p}\right) \left(q - 1 - 2\left\lfloor \frac{aq}{p} \right\rfloor\right).$$

Proof of Theorem 1

Before proceeding further, we recollect the fact that the class number $h(-p)$ is odd, which is our second lemma.

Lemma 2. *The class number $h(-p)$ of the imaginary quadratic field $\mathbb{Q}[\sqrt{-p}]$, $p \equiv 3 \bmod 4$, is odd.*

Proof. This fact is stated in [11].
Proceeding to the proof of the theorem, we find that when $p \equiv 1 \bmod q$, Lemma 1 may be written as

$$\left(q - \left(\frac{q}{p}\right)\right) h(-p)/2 = \sum_{k=1}^{(q-1)/2} (a_k - b_k) \left(\frac{q+1}{2} - k\right). \tag{0.2}$$

Noting that $a_k + b_k = \frac{p-1}{q}$, $k = 1, \ldots, (q-1)/2$, we may combine this with equation 0.2 to obtain

$$\left(\frac{q^2-1}{8}\right)\left(\frac{p-1}{2q}\right) - \frac{\left(q - \left(\frac{q}{p}\right)\right) h(-p)}{4} = \sum_{k=1}^{(q-1)/2} b_k \left(\frac{q+1}{2} - k\right). \tag{0.3}$$

The left hand side of equation 0.3 is always even.

Case $q \equiv 1 \bmod 8$. Then since $p \equiv 1 \bmod q$, by quadratic reciprocity, $\frac{\left(q - \left(\frac{q}{p}\right)\right)h(-p)}{4}$ is even, and considering $q \equiv 1, 9 \bmod 16$ shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-1}{2q}\right)$ is even.

Case $q \equiv 5 \bmod 8$. Then since $p \equiv 1 \bmod q$, by quadratic reciprocity, and by Lemma 2, $\frac{\left(q - \left(\frac{q}{p}\right)\right)h(-p)}{4}$ is odd. Considering $q \equiv 5, 13 \bmod 16$ shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-1}{2q}\right)$ is also odd.

Case $q \equiv 3 \bmod 8$. Then since $p \equiv 1 \bmod q$, by quadratic reciprocity, and by Lemma 2, $\frac{\left(q - \left(\frac{q}{p}\right)\right)h(-p)}{4}$ is odd. Considering $q \equiv 3, 11 \bmod 16$ shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-1}{2q}\right)$ is also odd.

Case $q \equiv 7 \bmod 8$. Then since $p \equiv 1 \bmod q$, by quadratic reciprocity, $\frac{\left(q - \left(\frac{q}{p}\right)\right)h(-p)}{4}$ is even. Considering $q \equiv 7, 15 \bmod 16$ shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-1}{2q}\right)$ is even.
Reducing equation 0.3 modulo 2, we obtain

$$\sum_{\substack{k=1 \\ (q+1)/2 - k \equiv 1 \bmod 2}}^{(q-1)/2} b_k \equiv 0 \bmod 2.$$

This proves Theorem 1.

Relevance for multiplicative orders of partial products and applications

In general, Wilson's theorem tells us that the product of all the products

$$\prod_{k=1}^{q} \Pi_k^{(q)}$$

is a quadratic non-residue for $p \equiv 3 \bmod 4$. Let q be an odd prime. Let $p \equiv 3 \bmod 4$ and $p \equiv 1 \bmod q$. Since $(p-1)/q$ is even, one has $\Pi_k^{(q)} \equiv \Pi_{q-k}^{(q)} \bmod p$, so $\Pi_{(q+1)/2}^{(q)}$ is a non-residue. It is thus interesting to ask which of the products $\Pi_k^{(q)}$, $k = 1, \ldots, (q-1)/2$ are quadratic residues, in which case they will have odd multiplicative order modulo p. We have shown that for p satisfying $p \equiv 3 \bmod 4$ and $p \equiv 1 \bmod q$, the product of the products $\prod_{\substack{k=1 \\ (q+1)/2-k \equiv 1 \bmod 2}}^{(q-1)/2} \Pi_k^{(q)}$ always has odd multiplicative order modulo p. Moreover, we may say the following.

Corollary. *Let q be an odd prime. Let $p \equiv 3 \bmod 4$ and $p \equiv 1 \bmod q$. Consider the set of products $\{\Pi_k^{(q)} : k = 1, \ldots, (q-1)/2, \text{ and } (q+1)/2 - k \equiv 1 \bmod 2\}$. The number of elements in this set that are quadratic non-residues is even. Thus, the number of elements in this set that have even multiplicative order modulo p is even.*

As examples of the corollary, we consider some cases for small q.
Case $q = 3$. Then $\Pi_1^{(3)}$ is a quadratic residue and has odd order modulo p.
Case $q = 5$. Then $\Pi_2^{(5)}$ is a quadratic residue and has odd order modulo p.
Case $q = 7$. Then $\Pi_1^{(7)}\Pi_3^{(7)}$ is a quadratic residue and has odd order modulo p. Either both $\Pi_1^{(7)}$, $\Pi_3^{(7)}$ are quadratic residues, or both are quadratic non-residues.
Case $q = 11$. Then $\Pi_1^{(11)}\Pi_3^{(11)}\Pi_5^{(11)}$ is a quadratic residue modulo p.

Further generalizations

Theorem 1 may be generalized to primes p such that $p \equiv 3 \bmod 4$ and $p \not\equiv 1 \bmod q$. In this case, we need a new generalization of the partial products $\Pi_k^{(q)}$. With this in mind, set

$$\Pi_k'^{(q)} = \prod_{j=\lfloor \frac{(k-1)p}{q} \rfloor + 1}^{\lfloor \frac{kp}{q} \rfloor} j,$$

$1 \leq k \leq q-1$,

$$\Pi_q'^{(q)} = \prod_{j=\lfloor \frac{(q-1)p}{q} \rfloor + 1}^{p-1} j.$$

We have that $\Pi_k'^{(q)} = \Pi_k^{(q)}$ when $p \equiv 1 \bmod q$. Let a_k', b_k', $1 \leq k \leq q-1$ denote respectively the number of quadratic residues and nonresidues in $\{\lfloor \frac{(k-1)p}{q} \rfloor + 1, \ldots, \lfloor \frac{kp}{q} \rfloor\}$, and a_q', b_q' denote respectively the number of quadratic residues and nonresidues in $\{\lfloor \frac{(q-1)p}{q} \rfloor + 1, \ldots, p-1\}$. These agree with the definition of a_k, b_k when $p \equiv 1 \bmod q$. Let us consider the case $p \equiv 2 \bmod q$. Then we have

Theorem 3. *Let q be an odd prime. Let p be a prime such that $p \equiv 3 \bmod 4$ and $p \equiv 2 \bmod q$. Then the element*

$$b = \prod_{\substack{k=1 \\ (q+1)/2-k \equiv 1 \bmod 2}}^{(q-1)/2} \Pi_k'^{(q)}$$

satisfies

$$\left(\frac{b}{p}\right) = \begin{cases} 1, \text{ if } q \equiv \pm 1 \bmod 16, \\ -1, \text{ if } q \equiv \pm 7 \bmod 16 \\ (-1)^{\frac{h(-p)+1}{2}}, \text{ if } q \equiv \pm 3 \bmod 16 \\ (-1)^{1+\frac{h(-p)+1}{2}}, \text{ if } q \equiv \pm 5 \bmod 16. \end{cases}$$

To verify this, we begin by checking that $a'_k + b'_k = \frac{p-2}{q}$, $1 \le k \le q$ and $k \ne (q+1)/2$, and $a'_k + b'_k = \frac{p-2}{q} + 1$ when $k = (q+1)/2$. Combining with Lemma 1 yields

$$\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right) - \frac{\left(q - \left(\frac{q}{p}\right)\right)h(-p)}{4} = \sum_{k=1}^{(q-1)/2} b'_k \left(\frac{q+1}{2} - k\right).$$

Case $q \equiv 1 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is even, and checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is even.
Case $q \equiv 9 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is even, and checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is odd.
Case $q \equiv 7 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is even, and checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is odd.
Case $q \equiv 15 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is even, and checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is even.
Case $q \equiv 5 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is a half-integer with numerator congruent to $3h(-q) \bmod 4$. Checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is a half-integer with numerator congruent to 3 mod 4.
Case $q \equiv 13 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is a half-integer with numerator congruent to $3h(-q) \bmod 4$. Checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is a half-integer with numerator congruent to 1 mod 4.
Case $q \equiv 3 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is a half-integer with numerator congruent to $h(-q) \bmod 4$. Checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is a half-integer with numerator congruent to 3 mod 4.
Case $q \equiv 11 \bmod 16$. Then $\frac{\left(q-\left(\frac{q}{p}\right)\right)h(-p)}{4}$ is a half-integer with numerator congruent to $h(-q) \bmod 4$. Checking q mod 32 shows that $\left(\frac{q^2-1}{8}\right)\left(\frac{p-2}{2q}\right)$ is a half-integer with numerator congruent to 1 mod 4.
Next is the case $p \equiv 3 \bmod q$.

Theorem 4. *Let $q > 3$ be an odd prime. Let p be a prime such that $p \equiv 3 \bmod 4$ and $p \equiv 3 \bmod q$. Then the element*

$$b = \prod_{\substack{k=1 \\ (q+1)/2-k \equiv 1 \bmod 2}}^{(q-1)/2} \Pi_k'^{(q)}$$

satisfies

$$\left(\frac{b}{p}\right) = \begin{cases} 1, & \text{if } q \equiv \pm 1 \bmod 12, \\ (-1)^{\frac{h(-p)+1}{2}}, & \text{if } q \equiv \pm 5 \bmod 12 \end{cases}$$

To verify this, we begin by checking that $a'_k + b'_k = \frac{p-3}{q}$, $1 \le k \le (q-1)/2$ and $k \ne \frac{q+2+\left(\left(\frac{q}{3}\right)-1\right)/2}{3}$, and $a'_k + b'_k = \frac{p-3}{q} + 1$ when $k = \frac{q+2+\left(\left(\frac{q}{3}\right)-1\right)/2}{3}$. Combining with Lemma 1 yields

$$\left(\frac{q^2-1}{8}\right)\left(\frac{p-3}{2q}\right) + \frac{q - \left(\frac{q}{3}\right)}{12} - \frac{\left(q - \left(\frac{q}{p}\right)\right)h(-p)}{4} = \sum_{k=1}^{(q-1)/2} b'_k \left(\frac{q+1}{2} - k\right),$$

the left hand side of the above being

$$\frac{\left(q - \left(\frac{q}{3}\right)\right)(1 - 3h(-p))}{12}$$

mod 2.

References

[1] H. Cohen, A Course in Computational Algebraic Number Theory, Springer Graduate Texts in Mathematics, No. 138, Berlin-Heidelberg 1993.
[2] J. B. Cosgrave and K. Dilcher, Extensions of the Gauss-Wilson Theorem, Integers: Electronic Journal of

Combinatorial Number Theory 8 (2008), A39.
[3] J. B. Cosgrave and K. Dilcher, Mod p^3 Analogues of Theorems of Gauss and Jacobi on Binomial Coefficients, Acta Arithmetica, Vol. 142, No. 2, 103–108, 2010.
[4] J. B. Cosgrave and K. Dilcher, The Multiplicative Order of Certain Gauss Factorials, International Journal of Number Theory, Vol. 7 (2011), 145–171.
[5] J. B. Cosgrave and K. Dilcher, An Introduction to Gauss Factorials, The American Mathematical Monthly, Vol. 118, No. 9, 812–829.
[6] J. B. Cosgrave and K. Dilcher, Sums of Reciprocals Modulo Composite Integers, Journal of Number Theory, 133, (2013), 3565–3577.
[7] J. B. Cosgrave and K. Dilcher, The Gauss-Wilson Theorem for Quarter-Intervals, Acta Mathematica Hungarica, 142, (2014), 199–230.
[8] J. B. Cosgrave and K. Dilcher, The Multiplicative Order of Certain Gauss Factorials II, Functiones et Approximatio Commentarii Mathematici.
[9] J. B. Cosgrave and K. Dilcher, A Role for Generalized Fermat Numbers, Mathematics of Computation, September 2015.
[10] H. Davenport, Multiplicative Number Theory, Second Edition, Graduate Texts in Mathematics, Vol. 74, Springer-Verlag, New York, 1980.
[11] A. Laradji, M. Mignotte, and N. Tzanakis, Elementary Trigonometric Sums Related to Quadratic Residues, Elemente Der Mathematik, Vol. 67, Issue 2, 2012, 51–60.
[12] S. G. Hahn and D. H. Lee, Some Congruences for Binomial Coefficients, Class Field Theory - Its Centenary and Prospect, Advanced Studies in Pure Mathemmatics, 30, Math. Soc. Japan (2001).
[13] S. G. Hahn and D. H. Lee, Gauss Sums and Binomial Coefficients, Journal of Number Theory, Volume 92, Issue 2, 2002, 257–271.
[14] L. J. Mordell, The Congruence $((p-1)/2)! \equiv \pm 1 (\mathrm{mod} p)$, American Mathematical Monthly 68 (1961), 145–146.

School of Computing,
National University of Singapore, Singapore 119077
Email: tchfoo@hotmail.com